

2015 HIV Fast Facts[‡]



Division of Public and Behavioral Health

Office of Public Health Informatics and Epidemiology

HIV/AIDS Surveillance Program



Brian Sandoval, Governor State of Nevada

Richard Whitley, MS, Director

Department of Health and Human Services

Cody L. Phinney, MPH, Administrator Division of Public and Behavioral Health

Leon Ravin, MD
Acting Chief Medical Officer

Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Fast Facts Nevada New HIV Diagnoses, 2015

| KE | | Total | | | Male | | | Female | |
|-----------------------------------|-----|-------|-------|-----|------|-------|----|--------|-------|
| | N | %^ | Rate* | n | %^ | Rate* | n | %^ | Rate* |
| County of Diagnosis | | | | | | | | | |
| Clark | 432 | 91% | 20.6 | 374 | 90% | 35.5 | 58 | 97% | 5.6 |
| Washoe | 38 | 8% | 8.6 | 36 | 9% | 16.2 | 2 | 3% | ~ |
| All Other Counties** | 7 | 1% | ~ | 7 | 2% | ~ | 0 | 0% | 0.0 |
| Total | 477 | 100% | 16.6 | 417 | 100% | 28.8 | 60 | 100% | 4.2 |
| Race/Ethnicity | | | | | | | | | |
| White, non-Hispanic | 164 | 34% | 10.7 | 144 | 35% | 18.6 | 20 | 33% | 2.6 |
| Black, non-Hispanic | 118 | 25% | 48.5 | 92 | 22% | 75.2 | 26 | 43% | 21.5 |
| Hispanic | 147 | 31% | 18.1 | 139 | 33% | 33.6 | 8 | 13% | ~ |
| Asian/Hawaiian/ Pacific Islander | 38 | 8% | 14.8 | 34 | 8% | 28.5 | 4 | 7% | ~ |
| American Indian/ Alaska Native | 4 | 1% | ~ | 2 | 0% | ~ | 2 | 3% | ~ |
| Multi-race/Other | 6 | 1% | NA | 6 | 1% | NA | 0 | 0% | NA |
| Total | 477 | 100% | 16.6 | 417 | 100% | 28.8 | 60 | 100% | 4.2 |
| Age Group at Diagnosis | | | | | | | | | |
| < 13 | 2 | 0% | ~ | 0 | 0% | 0.0 | 2 | 3% | ~ |
| 13 to 24 | 82 | 17% | 18.1 | 75 | 18% | 32.1 | 7 | 12% | ~ |
| 25 to 34 | 163 | 34% | 42.2 | 144 | 35% | 73.0 | 19 | 32% | 10.0 |
| 35 to 44 | 114 | 24% | 28.5 | 102 | 24% | 49.9 | 12 | 20% | 6.1 |
| 45 to 54 | 87 | 18% | 22.3 | 75 | 18% | 37.5 | 12 | 20% | 6.3 |
| 55 to 64 | 22 | 5% | 6.4 | 17 | 4% | 10.0 | 5 | 8% | ~ |
| 65 + | 7 | 1% | ~ | 4 | 1% | ~ | 3 | 5% | ~ |
| Total | 477 | 100% | 16.6 | 417 | 100% | 28.8 | 60 | 100% | 4.2 |
| Transmission Category | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 315 | 66% | NA | 315 | 76% | NA | 0 | 0% | NA |
| Injection drug use (IDU) | 18 | 4% | NA | 12 | 3% | NA | 6 | 10% | NA |
| MSM+IDU | 23 | 5% | NA | 23 | 6% | NA | 0 | 0% | NA |
| Heterosexual contact | 32 | 7% | NA | 12 | 3% | NA | 20 | 33% | NA |
| Perinatal exposure | 1 | 0% | NA | 0 | 0% | NA | 1 | 2% | NA |
| Transfusion/Hemophilia | 0 | 0% | NA | 0 | 0% | NA | 0 | 0% | NA |
| No identified risk (NIR) | 88 | 18% | NA | 55 | 13% | NA | 33 | 55% | NA |
| Total | 477 | 100% | 16.6 | 417 | 100% | 28.8 | 60 | 100% | 4.2 |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties

[~] Rates are suppressed due to unreliability based on the relative standard error for counts <12.

Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

 $^{^{\}Lambda} Percent's$ may not equal 100% due to rounding and unknown counts.



| YEE THE STATE OF T | | Total | | | Male | | | Female | ! |
|--|-----|-------|-------|-----|------|-------|----|--------|-------|
| | N | %^ | Rate* | n | %^ | Rate* | n | %^ | Rate* |
| County of Diagnosis | | | | | | | | | |
| Clark | 185 | 92% | 8.8 | 145 | 90% | 13.8 | 40 | 98% | 3.8 |
| Washoe | 11 | 5% | ~ | 10 | 6% | ~ | 1 | 2% | ~ |
| All Other Counties** | 6 | 3% | ~ | 6 | 4% | ~ | 0 | 0% | 0.0 |
| Total | 202 | 100% | 7.0 | 161 | 100% | 11.1 | 41 | 100% | 2.9 |
| Race/Ethnicity | | | | | | | | | |
| White, non-Hispanic | 79 | 39% | 5.2 | 67 | 42% | 8.7 | 12 | 29% | 1.6 |
| Black, non-Hispanic | 59 | 29% | 24.3 | 40 | 25% | 32.7 | 19 | 46% | 15.7 |
| Hispanic | 50 | 25% | 6.2 | 43 | 27% | 10.4 | 7 | 17% | ~ |
| Asian/Hawaiian/ Pacific Islander | 10 | 5% | 3.9 | 7 | 4% | ~ | 3 | 7% | ~ |
| American Indian/ Alaska Native | 1 | 0% | ~ | 1 | 1% | ~ | 0 | 0% | 0.0 |
| Multi-race/Other | 3 | 1% | NA | 3 | 2% | NA | 0 | 0% | NA |
| Total | 202 | 100% | 7.0 | 161 | 100% | 11.1 | 41 | 100% | 2.9 |
| Age Group at Diagnosis | | | | | | | | | |
| <13 | 0 | 0% | 0.0 | 0 | 0% | 0.0 | 0 | 0% | 0.0 |
| 13 to 24 | 15 | 7% | 3.3 | 10 | 6% | ~ | 5 | 12% | ~ |
| 25 to 34 | 53 | 26% | 13.7 | 49 | 30% | 24.8 | 4 | 10% | ~ |
| 35 to 44 | 59 | 29% | 14.7 | 48 | 30% | 23.5 | 11 | 27% | ~ |
| 45 to 54 | 53 | 26% | 13.6 | 39 | 24% | 19.5 | 14 | 34% | 7.3 |
| 55 to 64 | 13 | 6% | 3.8 | 9 | 6% | ~ | 4 | 10% | ~ |
| 65 + | 9 | 4% | ~ | 6 | 4% | ~ | 3 | 7% | ~ |
| Total | 202 | 100% | 7.0 | 161 | 100% | 11.1 | 41 | 100% | 2.9 |
| Transmission Category | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 118 | 58% | NA | 118 | 73% | NA | 0 | 0% | NA |
| Injection drug use (IDU) | 13 | 6% | NA | 7 | 4% | NA | 6 | 15% | NA |
| MSM+IDU | 8 | 4% | NA | 8 | 5% | NA | 0 | 0% | NA |
| Heterosexual contact | 18 | 9% | NA | 2 | 1% | NA | 16 | 39% | NA |
| Perinatal exposure | 0 | 0% | NA | 0 | 0% | NA | 0 | 0% | NA |
| Transfusion/Hemophilia | 0 | 0% | NA | 0 | 0% | NA | 0 | 0% | NA |
| No identified risk (NIR) | 45 | 22% | NA | 26 | 16% | NA | 19 | 46% | NA |
| Total | 202 | 100% | 7.0 | 161 | 100% | 11.1 | 41 | 100% | 2.9 |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

New stage 3 (AIDS) diagnoses are counted in eHARS surveillance statistics and include stage 3 (AIDS) and HIV cases diagnosed in Nevada, both living and deceased. The surveillance data excludes HIV cases diagnosed in other states but who currently live in Nevada. Stage 3 (AIDS) diagnoses and HIV diagnoses may duplicate case counts if the person was diagnosed with both stage 3 (AIDS) and HIV in the same year.

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

 $[\]sim$ Rates are suppressed due to unreliability based on the relative standard error for counts <12.

[§]Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

[^]Percent's may not equal 100% due to rounding and unknown counts.



Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Fast Facts Persons Living with HIV§ in Nevada, 2015

| XE SE | | Total | | 5.13 2 .11.18 | Male | | | Female | |
|-----------------------------------|--------|-------|---------|----------------------|------|---------|-------|--------|-------|
| | N | %^ | Rate* | n | %^ | Rate* | n | %^ | Rate* |
| County of Residence | | | | | | | | | |
| Clark | 8,753 | 86% | 417.6 | 7,357 | 86% | 699.0 | 1,396 | 87% | 133.8 |
| Washoe | 970 | 10% | 220.0 | 832 | 10% | 374.7 | 138 | 9% | 63.0 |
| All Other Counties** | 415 | 4% | 123.0 | 338 | 4% | 196.3 | 77 | 5% | 46.6 |
| Total | 10,138 | 100% | 352.7 | 8,527 | 100% | 589.4 | 1,611 | 100% | 112.9 |
| Race/Ethnicity | | | | | | | | | |
| White, non-Hispanic | 4,651 | 46% | 303.9 | 4,103 | 48% | 529.7 | 548 | 34% | 72.5 |
| Black, non-Hispanic | 2,547 | 25% | 1,048.0 | 1,820 | 21% | 1,487.0 | 727 | 45% | 601.9 |
| Hispanic | 2,349 | 23% | 289.6 | 2,095 | 25% | 505.7 | 254 | 16% | 64.0 |
| Asian/Hawaiian/ Pacific Islander | 368 | 4% | 143.4 | 318 | 4% | 266.6 | 50 | 3% | 36.4 |
| American Indian/Alaska Native | 82 | 1% | 252.2 | 63 | 1% | 387.5 | 19 | 1% | 116.9 |
| Multi-race/Other | 141 | 1% | NA | 128 | 2% | NA | 13 | 1% | NA |
| Total | 10,138 | 100% | 352.7 | 8,527 | 100% | 589.4 | 1,611 | 100% | 112.9 |
| Age Group at End of Year | | | | | | | | | |
| < 13 | 15 | 0% | 3.0 | 4 | 0% | ~ | 11 | 1% | ~ |
| 13 to 24 | 365 | 4% | 80.5 | 312 | 4% | 133.3 | 53 | 3% | 24.1 |
| 25 to 34 | 1,649 | 16% | 426.7 | 1,413 | 17% | 716.0 | 236 | 15% | 124.8 |
| 35 to 44 | 2,128 | 21% | 531.1 | 1,757 | 21% | 859.2 | 371 | 23% | 189.1 |
| 45 to 54 | 3,412 | 34% | 872.9 | 2,910 | 34% | 1,455.0 | 502 | 31% | 262.9 |
| 55 to 64 | 1,895 | 19% | 552.2 | 1,575 | 18% | 929.6 | 320 | 20% | 184.2 |
| 65 + | 616 | 6% | 154.4 | 507 | 6% | 274.0 | 109 | 7% | 50.9 |
| Missing | 58 | 1% | NA | 49 | 1% | NA | 9 | 1% | NA |
| Total | 10,138 | 100% | 352.7 | 8,527 | 100% | 589.4 | 1,611 | 100% | 112.9 |
| Transmission Category | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 6,484 | 64% | NA | 6,484 | 76% | NA | 0 | 0% | NA |
| Injection drug use (IDU) | 747 | 7% | NA | 498 | 6% | NA | 249 | 15% | NA |
| MSM+IDU | 665 | 7% | NA | 665 | 8% | NA | 0 | 0% | NA |
| Heterosexual contact | 1,271 | 13% | NA | 304 | 4% | NA | 967 | 60% | NA |
| Perinatal exposure | 75 | 1% | NA | 32 | 0% | NA | 43 | 3% | NA |
| Transfusion/Hemophilia | 11 | 0% | NA | 7 | 0% | NA | 4 | 0% | NA |
| No identified risk (NIR) | 885 | 9% | NA | 537 | 6% | NA | 348 | 22% | NA |
| Total | 10,138 | 100% | 352.7 | 8,527 | 100% | 589.4 | 1,611 | 100% | 112.9 |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

Persons living with HIV data include data on persons living in Nevada with HIV (not yet stage 3 (AIDS)) and stage 3 (AIDS) based on the current address listed in the HIV/AIDS Reporting System (eHARS). These persons may or may not have been diagnosed with HIV or stage 3 (AIDS) in Nevada.

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

 $^{^{\}sim}$ Rates are suppressed due to unreliability based on the relative standard error for counts <12.

[§]Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

[^]Percent's may not equal 100% due to rounding and unknown counts.

Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Fast Facts Clark County New HIV Diagnoses, 2015

| A KE | | Total | Cial | K County I | Male | gnoses, 2013 | | Female | |
|-----------------------------------|-----|-------|-------|------------|------|--------------|----|--------|-------|
| | N | %^ | Rate* | n | %^ | Rate* | n | %^ | Rate* |
| Race/Ethnicity | | | | | | | | | |
| White, non-Hispanic | 137 | 32% | 13.9 | 119 | 32% | 23.8 | 18 | 31% | 3.7 |
| Black, non-Hispanic | 115 | 27% | 50.5 | 89 | 24% | 78.6 | 26 | 45% | 22.7 |
| Hispanic | 137 | 32% | 21.1 | 129 | 34% | 39.1 | 8 | 14% | ~ |
| Asian/Hawaiian/ Pacific Islander | 33 | 8% | 14.9 | 29 | 8% | 28.2 | 4 | 7% | ~ |
| American Indian/ Alaska Native | 4 | 1% | ~ | 2 | 1% | ~ | 2 | 3% | ~ |
| Multi-race/Other | 6 | 1% | NA | 6 | 2% | NA | 0 | 0% | NA |
| Total | 432 | 100% | 20.6 | 374 | 100% | 35.5 | 58 | 100% | 5.6 |
| Age Group at Diagnosis | | | | | | | | | |
| < 13 | 2 | 0% | ~ | 0 | 0% | 0.0 | 2 | 3% | ~ |
| 13 to 24 | 75 | 17% | 22.9 | 68 | 18% | 40.4 | 7 | 12% | ~ |
| 25 to 34 | 148 | 34% | 52.2 | 130 | 35% | 90.1 | 18 | 31% | 12.9 |
| 35 to 44 | 106 | 25% | 34.1 | 94 | 25% | 59.4 | 12 | 21% | 7.9 |
| 45 to 54 | 78 | 18% | 27.0 | 66 | 18% | 44.7 | 12 | 21% | 8.5 |
| 55 to 64 | 16 | 4% | 6.7 | 12 | 3% | 10.2 | 4 | 7% | ~ |
| 65 + | 7 | 2% | ~ | 4 | 1% | ~ | 3 | 5% | ~ |
| Total | 432 | 100% | 20.6 | 374 | 100% | 35.5 | 58 | 100% | 5.6 |
| Transmission Category | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 288 | 67% | NA | 288 | 77% | NA | 0 | 0% | NA |
| Injection drug use (IDU) | 14 | 3% | NA | 8 | 2% | NA | 6 | 10% | NA |
| MSM+IDU | 19 | 4% | NA | 19 | 5% | NA | 0 | 0% | NA |
| Heterosexual contact | 31 | 7% | NA | 12 | 3% | NA | 19 | 33% | NA |
| Perinatal exposure | 1 | 0% | NA | 0 | 0% | NA | 1 | 2% | NA |
| Transfusion/Hemophilia | 0 | 0% | NA | 0 | 0% | NA | 0 | 0% | NA |
| No identified risk (NIR) | 79 | 18% | NA | 47 | 13% | NA | 32 | 55% | NA |
| Total | 432 | 100% | 20.6 | 374 | 100% | 35.5 | 58 | 100% | 5.6 |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

[~] Rates are suppressed due to unreliability based on the relative standard error for counts <12.

[§]Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

[^]Percent's may not equal 100% due to rounding and unknown counts.

Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Fast Facts Clark County New Stage 3 (AIDS) Diagnoses, 2015

| | | | Clark Col | unty New 3 | Stage 3 (AID: | S) Diagnoses, | 2015 | | |
|---|-----|-------|-----------|------------|---------------|---------------|------|------------|-------|
| A DENADOR | | Total | | | Male | | | Female | |
| | N | %^ | Rate* | n | %^ | Rate* | n | % ^ | Rate* |
| Race/Ethnicity | | | | | | | | | |
| White, non-Hispanic | 66 | 36% | 6.7 | 55 | 38% | 11.0 | 11 | 28% | 2.3 |
| Black, non-Hispanic | 58 | 31% | 25.5 | 39 | 27% | 34.4 | 19 | 48% | 16.6 |
| Hispanic | 49 | 26% | 7.5 | 42 | 29% | 12.7 | 7 | 18% | ~ |
| Asian/Hawaiian/ Pacific Islander | 8 | 4% | ~ | 5 | 3% | ~ | 3 | 8% | ~ |
| American Indian/ Alaska Native | 1 | 1% | ~ | 1 | 1% | ~ | 0 | 0% | 0.0 |
| Multi-race/Other | 3 | 2% | NA | 3 | 2% | NA | 0 | 0% | NA |
| Total | 185 | 100% | 8.8 | 145 | 100% | 13.8 | 40 | 100% | 3.8 |
| Age Group at Diagnosis | | | | | | | | | |
| < 13 | 0 | 0% | 0.0 | 0 | 0% | 0.0 | 0 | 0% | 0.0 |
| 13 to 24 | 13 | 7% | 4.0 | 8 | 6% | ~ | 5 | 13% | ~ |
| 25 to 34 | 51 | 28% | 18.0 | 47 | 32% | 32.6 | 4 | 10% | ~ |
| 35 to 44 | 56 | 30% | 18.0 | 45 | 31% | 28.4 | 11 | 28% | ~ |
| 45 to 54 | 48 | 26% | 16.6 | 34 | 23% | 23.0 | 14 | 35% | 9.9 |
| 55 to 64 | 9 | 5% | ~ | 6 | 4% | ~ | 3 | 8% | ~ |
| 65 + | 8 | 4% | ~ | 5 | 3% | ~ | 3 | 8% | ~ |
| Total | 185 | 100% | 8.8 | 145 | 100% | 13.8 | 40 | 100% | 3.8 |
| Transmission Category | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 108 | 58% | NA | 108 | 74% | NA | 0 | 0% | NA |
| Injection drug use (IDU) | 11 | 6% | NA | 6 | 4% | NA | 5 | 13% | NA |
| MSM+IDU | 6 | 3% | NA | 6 | 4% | NA | 0 | 0% | NA |
| Heterosexual contact | 18 | 10% | NA | 2 | 1% | NA | 16 | 40% | NA |
| Perinatal exposure | 0 | 0% | NA | 0 | 0% | NA | 0 | 0% | NA |
| Transfusion/ Hemophilia | 0 | 0% | NA | 0 | 0% | NA | 0 | 0% | NA |
| No identified risk (NIR) | 42 | 23% | NA | 23 | 16% | NA | 19 | 48% | NA |
| Total Source: Division of Bublic and Robavioral Health | 185 | 100% | 8.8 | 145 | 100% | 13.8 | 40 | 100% | 3.8 |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

New stage 3 (AIDS) diagnoses are counted in eHARS surveillance statistics and include stage 3 (AIDS) and HIV cases diagnosed in Nevada, both living and deceased. The surveillance data excludes HIV cases diagnosed in other states but who currently live in Nevada. Stage 3 (AIDS) diagnoses and HIV diagnoses may duplicate case counts if the person was diagnosed with both stage 3 (AIDS) and HIV in the same year.

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

 $[\]sim$ Rates are suppressed due to unreliability based on the relative standard error for counts <12.

[§]Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

[^]Percent's may not equal 100% due to rounding and unknown counts.



| X. S. | | Total | | | Male | | | Female | |
|---|-------|-------|---------|-------|------------|---------|-------|--------|-------|
| | N | %^ | Rate* | n | % ^ | Rate* | n | %^ | Rate* |
| Race/Ethnicity | | | | | | | | | |
| White, non-Hispanic | 3,744 | 43% | 380.3 | 3,333 | 45% | 666.7 | 411 | 29% | 84.8 |
| Black, non-Hispanic | 2,392 | 27% | 1,050.0 | 1,694 | 23% | 1,496.0 | 698 | 50% | 609.6 |
| Hispanic | 2,097 | 24% | 322.9 | 1,880 | 26% | 570.5 | 217 | 16% | 67.9 |
| Asian/Hawaiian/ Pacific Islander | 336 | 4% | 152.2 | 293 | 4% | 285.0 | 43 | 3% | 36.5 |
| American Indian/ Alaska Native | 61 | 1% | 446.6 | 46 | 1% | 660.1 | 15 | 1% | 224.2 |
| Multi-race/Other | 123 | 1% | NA | 111 | 2% | NA | 12 | 1% | NA |
| Total | 8,753 | 100% | 417.6 | 7,357 | 100% | 699.0 | 1,396 | 100% | 133.8 |
| Age Group at End of Year | | | | | | | | | |
| < 13 | 13 | 0% | 3.5 | 4 | 0% | ~ | 9 | 1% | ٧ |
| 13 to 24 | 324 | 4% | 98.8 | 275 | 4% | 163.5 | 49 | 4% | 30.7 |
| 25 to 34 | 1,478 | 17% | 521.0 | 1,273 | 17% | 882.5 | 205 | 15% | 147.0 |
| 35 to 44 | 1,881 | 21% | 605.6 | 1,549 | 21% | 978.6 | 332 | 24% | 217.9 |
| 45 to 54 | 2,890 | 33% | 999.2 | 2,470 | 34% | 1,672.0 | 420 | 30% | 296.8 |
| 55 to 64 | 1,599 | 18% | 670.9 | 1,323 | 18% | 1,128.0 | 276 | 20% | 228.0 |
| 65 + | 511 | 6% | 188.2 | 414 | 6% | 331.0 | 97 | 7% | 66.3 |
| Missing | 57 | 1% | NA | 49 | 1% | NA | 8 | 1% | NA |
| Total | 8,753 | 100% | 417.6 | 7,357 | 100% | 699.0 | 1,396 | 100% | 133.8 |
| Transmission Category | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 5,737 | 66% | NA | 5,737 | 78% | NA | 0 | 0% | NA |
| Injection drug use (IDU) | 589 | 7% | NA | 391 | 5% | NA | 198 | 14% | NA |
| MSM+IDU | 534 | 6% | NA | 534 | 7% | NA | 0 | 0% | NA |
| Heterosexual contact | 1122 | 13% | NA | 257 | 3% | NA | 865 | 62% | NA |
| Perinatal exposure | 67 | 1% | NA | 30 | 0% | NA | 37 | 3% | NA |
| Transfusion/Hemophilia | 9 | 0% | NA | 7 | 0% | NA | 2 | 0% | NA |
| No identified risk (NIR) | 695 | 8% | NA | 401 | 5% | NA | 294 | 21% | NA |
| Total | 8,753 | 100% | 417.6 | 7,357 | 100% | 699.0 | 1,396 | 100% | 133.8 |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

Persons living with HIV data include data on persons living in Nevada with HIV (not yet stage 3 (AIDS)) and stage 3 (AIDS) based on the current address listed in the HIV/AIDS Reporting System (eHARS). These persons may or may not have been diagnosed with HIV or stage 3 (AIDS) in Nevada.

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

 $[\]sim$ Rates are suppressed due to unreliability based on the relative standard error for counts <12.

[§]Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

[^]Percent's may not equal 100% due to rounding and unknown counts.

Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Fast Facts Washoe County New HIV Diagnoses and Stage 3 (AIDS) Diagnoses, 2015 New HIV Diagnoses New Stage 3 (AIDS) Diagnoses

| X KE | | New HIV Diagnose | | | ew Stage 3 (AIDS) Dia | |
|---------------------------------|----|------------------|-------|----|-----------------------|-------|
| | N | %^ | Rate* | N | %^ | Rate* |
| Sex at Birth | | | | | | |
| Male | 36 | 95% | 16.2 | 10 | 91% | ~ |
| Female | 2 | 5% | ~ | 1 | 9% | ~ |
| Total | 38 | 100% | 8.6 | 11 | 100% | ~ |
| Race/Ethnicity | | | | | | |
| White, non-Hispanic | 22 | 58% | 7.7 | 8 | 73% | ~ |
| Black, non-Hispanic | 3 | 8% | ~ | 1 | 9% | ~ |
| Hispanic | 8 | 21% | ~ | 0 | 0% | 0.0 |
| Asian/Hawaiian/Pacific Islander | 5 | 13% | ~ | 2 | 18% | ~ |
| American Indian/Alaska Native | 0 | 0% | 0.0 | 0 | 0% | 0.0 |
| Multi-race/Other | 0 | 0% | NA | 0 | 0% | NA |
| Total | 38 | 100% | 8.6 | 11 | 100% | ~ |
| Age Group at Diagnosis | | | | | | |
| <13 | 0 | 0% | 0.0 | 0 | 0% | 0.0 |
| 13 to 24 | 6 | 16% | ~ | 2 | 18% | ~ |
| 25 to 34 | 13 | 34% | 20.5 | 2 | 18% | ~ |
| 35 to 44 | 6 | 16% | ~ | 1 | 9% | ~ |
| 45 to 54 | 7 | 18% | ~ | 2 | 18% | ~ |
| 55 to 64 | 6 | 16% | ~ | 4 | 36% | ~ |
| 65 + | 0 | 0% | 0.0 | 0 | 0% | 0.0 |
| Total | 38 | 100% | 8.6 | 11 | 100% | ~ |
| Transmission Category | | | | | | |
| Males | | | | | | |
| Male-to-male sexual contact | 22 | C 40/ | NA | 7 | 700/ | NA |
| (MSM) | 23 | 64% | NA | 7 | 70% | NA |
| Injection drug use (IDU) | 4 | 11% | NA | 1 | 10% | NA |
| MSM+IDU | 3 | 8% | NA | 1 | 10% | NA |
| Heterosexual contact | 0 | 0% | NA | 0 | 0% | NA |
| Perinatal Exposure | 0 | 0% | NA | 0 | 0% | NA |
| Transfusion/Hemophilia | 0 | 0% | NA | 0 | 0% | NA |
| No identified risk (NIR) | 6 | 17% | NA | 1 | 10% | NA |
| Subtotal | 36 | 100% | 16.2 | 10 | 100% | ~ |
| Females | | | | | | |
| IDU | 0 | 0% | NA | 1 | 100% | NA |
| Heterosexual contact | 1 | 50% | NA | 0 | 0% | NA |
| Perinatal Exposure | 0 | 0% | NA | 0 | 0% | NA |
| Transfusion/Hemophilia | 0 | 0% | NA | 0 | 0% | NA |
| No identified risk (NIR) | 1 | 50% | NA | 0 | 0% | NA |
| Subtotal | 2 | 100% | ~ | 1 | 100% | ~ |
| Total | 38 | 100% | 8.6 | 11 | 100% | ~ |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

[~] Rates are suppressed due to unreliability based on the relative standard error for counts <12.

 $[$]Persons \ Living \ with \ HIV \ indicate \ any \ person \ regardless \ of \ HIV \ staging, \ including \ HIV \ stage \ 3 \ (AIDS), \ living \ in \ Nevada \ in \ 2015.$

[^]Percent's may not equal 100% due to rounding and unknown counts.



Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Fast Facts Persons Living with HIV§ in Washoe County, 2015

| | | Total | | | Male | 7. | | Female | |
|-----------------------------------|-----|--------|--------|-----|--------|--------|-----|--------|-------|
| | N | %^ | Rate* | n | %^ | Rate* | n | %^ | Rate* |
| Race/Ethnicity | | | | | | | | | |
| White, non-Hispanic | 626 | 65% | 218.2 | 544 | 65% | 376.6 | 82 | 59% | 57.6 |
| Black, non-Hispanic | 110 | 11% | 1009.0 | 90 | 11% | 1503.0 | 20 | 15% | 407.0 |
| Hispanic | 179 | 19% | 168.5 | 152 | 18% | 279.6 | 27 | 20% | 52.0 |
| Asian/Hawaiian/Pacific Islander | 28 | 3% | 94.4 | 23 | 3% | 167.3 | 5 | 4% | ~ |
| American Indian/Alaska Native | 14 | 1% | 193.9 | 11 | 1% | ~ | 3 | 2% | ~ |
| Multi-race/Other/Unknown | 13 | 1% | 2.9 | 12 | 1% | ~ | 1 | 1% | ~ |
| Total | 970 | 100% | 220.0 | 832 | 100% | 374.7 | 138 | 100% | 63.0 |
| Age Group at End of Year | | | | | | | | | |
| < 13 | 0 | 0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0% | 0.0 |
| 13 to 24 | 34 | 4% | 48.3 | 31 | 3.7% | 85.1 | 3 | 2% | ~ |
| 25 to 34 | 133 | 14% | 209.5 | 109 | 13.1% | 336.9 | 24 | 17% | 77.1 |
| 35 to 44 | 176 | 18% | 322.8 | 151 | 18.1% | 545.6 | 25 | 18% | 93.1 |
| 45 to 54 | 358 | 37% | 623.7 | 303 | 36.4% | 1040.0 | 55 | 40% | 194.7 |
| 55 to 64 | 204 | 21% | 359.1 | 179 | 21.5% | 636.3 | 25 | 18% | 87.2 |
| 65 + | 65 | 7% | 103.4 | 59 | 7.1% | 200.4 | 6 | 4% | ~ |
| Missing | 0 | 0% | NA | 0 | 0.0% | NA | 0 | 0% | NA |
| Total | 970 | 100.0% | 220.0 | 832 | 100.0% | 374.7 | 138 | 100% | 63.0 |
| Transmission Category | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 560 | 58% | NA | 560 | 67% | NA | 0 | 0% | NA |
| Injection drug use (IDU) | 90 | 9% | NA | 62 | 8% | NA | 28 | 20% | NA |
| MSM+IDU | 91 | 9% | NA | 91 | 11% | NA | 0 | 0% | NA |
| Heterosexual contact | 96 | 10% | NA | 25 | 3% | NA | 71 | 51% | NA |
| Perinatal Exposure | 4 | 0 % | NA | 1 | 0% | NA | 3 | 2% | NA |
| Transfusion/Hemophilia | 0 | 0% | NA | 0 | 0% | NA | 0 | 0% | NA |
| No identified risk (NIR) | 129 | 13 % | NA | 93 | 11% | NA | 36 | 26% | NA |
| Total | 970 | 100% | 220.0 | 832 | 100% | 374.7 | 138 | 100% | 63.0 |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

Persons living with HIV data include data on persons living in Nevada with HIV (not yet stage 3 (AIDS)) and stage 3 (AIDS) based on the current address listed in the HIV/AIDS Reporting System (eHARS). These persons may or may not have been diagnosed with HIV or stage 3 (AIDS) in Nevada.

^{*} Rates per 100,000 population were calculated using 2015 population projections from the Nevada State Demographer vintage 2015 data.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

[~] Rates are suppressed due to unreliability based on the relative standard error for counts <12.

[§]Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

[^]Percent's may not equal 100% due to rounding and unknown counts.

Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Fast Facts Annual Percent Change: New HIV Diagnoses and Persons Living with HIV in Nevada, 2014-2015



| A STATE OF THE STA | | | New HI | V Diagnos | ses | | All | Persons Liv | ing with H | ΗV [§] |
|--|-----|------|--------|-----------|-----------------------|-------|------|-------------|------------|-----------------------|
| × | 20 |)14 | 20 | 015 | Annual % Change: | 20: | 14 | 201 | 15 | Annual % Change: |
| | N | %^ | N | %^ | % Change ¹ | N | %^ | N | %^ | % Change ¹ |
| County | | | | | | | | | | |
| Clark | 375 | 89% | 432 | 91% | 15% | 8,449 | 86% | 8,753 | 86% | 4% |
| Washoe | 37 | 9% | 38 | 8% | 3% | 956 | 10% | 970 | 10% | 1% |
| All Other Counties** | 8 | 2% | 7 | 1% | # | 426 | 4% | 415 | 4% | -3% |
| Total | 420 | 100% | 477 | 100% | 14% | 9,831 | 100% | 10,138 | 100% | 3% |
| Sex at Birth | | | | | | | | | | |
| Male | 364 | 87% | 417 | 87% | 15% | 8,282 | 84% | 8,527 | 84% | 3% |
| Female | 56 | 13% | 60 | 13% | 7% | 1,549 | 16% | 1,611 | 16% | 4% |
| Total | 420 | 100% | 477 | 100% | 14% | 9,831 | 100% | 10,138 | 100% | 3% |
| Race/Ethnicity | | | | | | | | | | |
| White, non-Hispanic | 155 | 37% | 164 | 34% | 6% | 4,652 | 47% | 4,651 | 46% | 0% |
| Black, non-Hispanic | 101 | 24% | 118 | 25% | 17% | 2,432 | 25% | 2,547 | 25% | 5% |
| Hispanic | 128 | 30% | 147 | 31% | 15% | 2,208 | 22% | 2,349 | 23% | 6% |
| Asian/Hawaiian/Pacific Islander | 23 | 5% | 38 | 8% | 65% | 330 | 3% | 368 | 4% | 12% |
| American Indian/Alaska Native | 3 | 1% | 4 | 1% | # | 80 | 1% | 82 | 1% | 3% |
| Multi-race/Other | 10 | 2% | 6 | 1% | # | 129 | 1% | 141 | 1% | 9% |
| Total | 420 | 100% | 477 | 100% | 14% | 9,831 | 100% | 10,138 | 100% | 3% |
| Age Group* | | | | | | | | | | |
| < 13 | 2 | 0% | 2 | 0% | # | 11 | 0% | 15 | 0% | 36% |
| 13 to 24 | 97 | 23% | 82 | 17% | -15% | 363 | 4% | 365 | 4% | 1% |
| 25 to 34 | 145 | 35% | 163 | 34% | 12% | 1,561 | 16% | 1,649 | 16% | 6% |
| 35 to 44 | 78 | 19% | 114 | 24% | 46% | 2,170 | 22% | 2,128 | 21% | -2% |
| 45 to 54 | 73 | 17% | 87 | 18% | 19% | 3,371 | 34% | 3,412 | 34% | 1% |
| 55 to 64 | 22 | 5% | 22 | 5% | 0% | 1,754 | 18% | 1,895 | 19% | 8% |
| 65 + | 3 | 1% | 7 | 1% | # | 543 | 6% | 616 | 6% | 13% |
| Missing | 0 | 0% | 0 | 0% | NA | 58 | 1% | 58 | 1% | 0% |
| Total | 420 | 100% | 477 | 100% | 14% | 9,831 | 100% | 10,138 | 100% | 3% |
| Transmission Category | | | | | | | | | | |
| Males | | | | | | | | | | |
| Male-to-male sexual contact (MSM) | 273 | 75% | 315 | 76% | 15% | 6,297 | 76% | 6,484 | 76% | 3% |
| IDU | 12 | 3% | 12 | 3% | 0% | 498 | 6% | 498 | 6% | 0% |
| MSM+IDU | 25 | 7% | 23 | 6% | -8% | 639 | 8% | 665 | 8% | 4% |
| Heterosexual contact | 11 | 3% | 12 | 3% | # | 294 | 4% | 304 | 4% | 3% |
| Perinatal Exposure | 0 | 0% | 0 | 0% | NA | 33 | 0% | 32 | 0% | -3% |
| Transfusion/Hemophilia | 0 | 0% | 0 | 0% | NA | 7 | 0% | 7 | 0% | NA" |
| No identified risk (NIR) | 43 | 12% | 55 | 13% | 28% | 514 | 6% | 537 | 6% | 4% |
| Subtotal | 364 | 100% | 417 | 100% | 15% | 8,282 | 100% | 8,527 | 100% | 3% |
| Females | | | | | | | | | | |
| IDU | 5 | 9% | 6 | 10% | + | 249 | 16% | 249 | 15% | 0% |
| Heterosexual contact | 20 | 36% | 20 | 33% | 0% | 936 | 60% | 967 | 60% | 3% |
| Perinatal Exposure | 2 | 4% | 1 | 2% | # | 40 | 3% | 43 | 3% | 8% |
| Transfusion/Hemophilia | 0 | 0% | 0 | 0% | NA | 3 | 0% | 4 | 0% | NA" |
| No identified risk (NIR) | 29 | 52% | 33 | 55% | 14% | 321 | 21% | 348 | 22% | 8% |
| Subtotal | 56 | 100% | 60 | 100% | 7% | 1,549 | 100% | 1,611 | 100% | 4% |
| Total | 420 | 100% | 477 | 100% | 14% | 9,831 | 100% | 10,138 | 100% | 3% |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

Persons living HIV data includes data on persons living in Nevada with HIV (not yet stage 3 (AIDS)) and stage 3 (AIDS) based on the current address listed in the HIV/AIDS Reporting System (eHARS). These persons may or may not have been diagnosed with HIV or stage 3 (AIDS) in Nevada.

Annual % Change is the percentage increase or decrease relative to the number of cases in 2014. Take caution when interpreting percent changes calculated from small case counts because they can be susceptible to high variability and are likely to be misleading.
"Percent change for small counts are not shown as they are susceptible to high variability and are misleading.

^{*} For New HIV diagnoses, age refers to age at HIV infection diagnosis. For persons living with HIV, age refers to age at the end of the report year.

^{**}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humbolat, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

§Persons Living with HIV indicate any person regardless of HIV staging, including HIV stage 3 (AIDS), living in Nevada in 2015.

[^] Percent's may not equal 100% due to rounding and unknown counts.

| Division of Public and Behavioral Health- HIV/AIDS Surveillance Program |
|--|
| Facility of Diagnosis for New HIV Diagnoses, Nevada HIV Fast Facts, 2015 |

| A CONTRACTOR OF THE CONTRACTOR | Nev | rada | Clark C | County | Washo | e County | All Other Counties* | |
|--|-----|------|---------|------------|-------|------------|------------------------|------|
| | N | %^ | n | % ^ | n | % ^ | n | %^ |
| Facility of Diagnosis | | | | | | | | |
| HIV Counseling and Testing Site | 124 | 26% | 115 | 27% | 9 | 24% | 0 | 0% |
| Private Physician's Office | 147 | 31% | 141 | 33% | 4 | 11% | 2 | 29% |
| Inpatient Facility/Hospital | 78 | 16% | 70 | 16% | 7 | 18% | 1 | 14% |
| Outpatient Facility (unspecified) | 24 | 5% | 14 | 3% | 9 | 24% | 1 | 14% |
| Adult HIV Clinic | 7 | 1% | 3 | 1% | 3 | 8% | 1 | 14% |
| Correctional Facility | 24 | 5% | 23 | 5% | 0 | 0% | 1 | 14% |
| STD Clinic | 42 | 9% | 41 | 9% | 0 | 0% | 1 | 14% |
| Blood Bank or Plasma Center | 18 | 4% | 14 | 3% | 4 | 11% | 0 | 0% |
| Emergency Room | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| Tuberculosis Clinic | 3 | 1% | 3 | 1% | 0 | 0% | 0 | 0% |
| Obstetrics and Gynecology Clinic | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% |
| Facility/Other/Unknown | 10 | 2% | 8 | 2% | 2 | 5% | 0 | 0% |
| Total | 477 | 100% | 432 | 100% | 38 | 100% | 7 | 100% |

| SI C NEVADA | | Division of Public and Behavioral Health- HIV/AIDS Surveillance Program Facility of Diagnosis for New Stage 3 (AIDS) Diagnoses, Nevada HIV Fast Facts, 2015 | | | | | | | | | |
|-----------------------------------|-----|---|---------|--------|-------|----------|------------------------|------|--|--|--|
| A KE | Nev | ada | Clark C | County | Washo | e County | All Other Counties* | | | | |
| | N | %^ | n | %^ | n | %^ | n | %^ | | | |
| Facility of Diagnosis | | | | | | | | | | | |
| HIV Counseling and Testing Site | 28 | 13% | 28 | 15% | 0 | 0% | 0 | 0% | | | |
| Private Physician's Office | 49 | 24% | 45 | 24% | 2 | 18% | 2 | 33% | | | |
| Inpatient Facility/Hospital | 88 | 44% | 82 | 45% | 3 | 27% | 3 | 50% | | | |
| Outpatient Facility (unspecified) | 1 | 0% | 0 | 0% | 1 | 9% | 0 | 0% | | | |
| Adult HIV Clinic | 15 | 7% | 10 | 5% | 5 | 45% | 0 | 0% | | | |
| Correctional Facility | 6 | 3% | 5 | 3% | 0 | 0% | 1 | 17% | | | |
| STD Clinic | 15 | 7% | 15 | 8% | 0 | 0% | 0 | 0% | | | |
| Blood Bank or Plasma Center | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | |
| Emergency Room | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | |
| Tuberculosis Clinic | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | |
| Obstetrics and Gynecology Clinic | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | |
| Facility/Other/Unknown | 0 | 0% | 0 | 0% | 0 | 0% | 0 | 0% | | | |
| Total | 202 | 100% | 185 | 100% | 11 | 100% | 6 | 100% | | | |

Source: Division of Public and Behavioral Health, HIV/AIDS Reporting System (eHARS), (March 2016)

^{*}All other counties include Carson City, Churchill, Douglas, Elko, Esmeralda, Eureka, Humboldt, Lander, Lincoln, Lyon, Mineral, Nye, Pershing, Storey, and White Pine Counties.

[^]Percent's may not equal 100% due to rounding and unknown counts.

For more information, contact:

Danika Williams, MPH
HIV Surveillance Coordinator
Office of Public Health Informatics and Epidemiology
Division of Public and Behavioral Health
4126 Technology Way Ste 201
Carson City, NV 89706
Phone: (775)684-2219
dmwilliams@health.nv.gov

James Kuzhippala, MPH
HIV/STD Epidemiologist
Office of Public Health Informatics and Epidemiology
Division of Public and Behavioral Health
4126 Technology Way Ste 201
Carson City, NV 89706
Phone: (775)684-5979
jkuzhippala@health.nv.gov

Theron Huntamer
HIV/AIDS Epidemiology Capacity Coordinator
Office of Public Health Informatics and Epidemiology
Division of Public and Behavioral Health
4126 Technology Way Ste 201
Carson City, NV 89706
Phone: (775)684-4152
thuntamer@health.nv.gov

Recommended Citation:

Office of Public Health Informatics and Epidemiology. Division of Public and Behavioral Health. *2015 HIV Fast Facts*. Carson City, Nevada. e1.0. June 2016.

Report Changes:

*"HIV" was previously referred to as "HIV/AIDS;" "Stage 3 (AIDS)" was previously referred to as "AIDS." The change in reference is due to a change in case definition, in which a staging system is used where AIDS is now end stage HIV (Stage 3) and HIV refers to all stages, including AIDS. More information can be found here:

http://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-us.pdf http://www.cdc.gov/mmwr/preview/mmwrhtml/rr6303a1.htm

The authors would like to recognize the Centers for Disease Control and Prevention (CDC) for their support through Grant Number 5U62PS004024 for the HIV/AIDS Surveillance Program. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC.