Influenza Weekly Report

2012 Week 51 (December 16 - 22) through 2013 Week 51 (December 15-21)

Department of Health and Human Services
Division of Public and Behavioral Health
Office of Public Health Informatics and Epidemiology



Brian Sandoval Governor State of Nevada

Michael J Willden Director Department of Health and Human Services

Richard Whitley, MS Administrator Division of Public and Behavioral Health

Tracey D Green, MD Chief Medical Officer Division of Public and Behavioral Health

Data for the graphs and tables on the following pages are provisional and may be updated as additional information becomes available.


Purpose

The purpose of this report is to provide an overview of and statistics for the influenza season in Nevada for the local public health authorities, sentinel providers and the public.

Influenza-Like Illness Network Surveillance (ILINet)

Respiratory specimens tested for influenza by the World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NRVESS) collaborating laboratories by sub-type. There were 13,548 specimens collected nationally during week 51 that were tested for influenza; of these 3,817 tested positive or the percent positive was 15.7%.


Figure 1



Source of Data: CDC: FluView Weekly Report.

Of the 62 specimens tested for influenza at both the Nevada State Public Health Laboratory and Southern Nevada Public Health Laboratory for sentinel providers, 35 were positive for influenza during week 51 or 56.5%.

Figure 2



Source of Data: CDC: ILINet.

Nevada State Public Health Laboratory (NSPHL) has tested 97 specimens this season with 72 positive from sentinel providers (74.2% positive). Southern Nevada Public Health Laboratory (SNPHL) has reported 8 positive influenza specimens through the Pediatric Early Warning Sentinel Surveillance (PEWSS). Nationally, there have been 90,428 specimens sent to the WHO and NERVSS laboratories with 11,391 positive or 12.6%. The national numbers in table 1 are reflected in figure 1. The state of Nevada data in table 1 is reflected in figure 2.


Table 1

ILINet Surveillance: Influenza Specimens Tested State and Nationally										
	NSPHL	SNPHL	State of Nevada (Week 51)		State of Nevada (Season)		National (Week 51)		National (Season)	
			#	%	#	%	#	%	#	%
Specimens Tested	97	115	62		212		13,548		90,428	
Positives to Influenza	72	8	35	56.5	80	37.7	3,817	28.2	11,391	12.6
<u>Influenza A:</u>	72	8	35	100.0	80	100.0	3,752	98.3	10,896	95.7
A(2009 H1N1)	71	8	35	100.0	79	98.8	2,111	56.3	5,998	55.0
A(Sub-typing not performed)	0	0	0	0.0	0	0.0	1,591	42.4	4,572	42.0
A(H3)	1	0	0	0.0	1	1.3	50	1.3	326	3.0
<u>Influenza B:</u>	0	0	0	0.0	0	0.0	65	1.7	494	4.3

Source of Data: CDC: FluView Report and CDC: ILINet.

For week 51, Nevada reported regional activity to CDC, along with 23 states and territories (Arkansas, Colorado, Connecticut, Florida, Georgia, Guam, Idaho, Illinois, Indiana, Iowa, Kentucky, Maine, Michigan, Minnesota, Mississippi, Nebraska, New Jersey, Oklahoma, Oregon, South Carolina, Tennessee, Utah, and Washington). Activity level¹ is derived from data analyzed from Influenza-like Illness (ILI) surveillance (laboratory and sentinel data), and data reported to the state through NBS/NETSS.

Figure 3




Source of Maps: CDC: FluView Report.

Influenza-like Illness (ILI) Network Surveillance has each sentinel providers report the number of patients that meet the ILI case definition² and number of patients that visit the provider weekly, which increased from 16,512 (week 50) to 17,453 (week 51). The "percentage of visits" is the number of ILI patients divided by the total number of patient visit per week. Nevada's ILI percentage of visits to providers increased to 2.6% during week 51, and is below the state baseline of 2.3%. Region 9 increased in ILI to 3.0% from 2.6%, and includes the following states/territories: Arizona, California, Guam, Hawaii, and Nevada. The nation increased to 3.2% from 2.4% during week 51.

^{1:} Activity level: Appendix Table 4.

^{2:} ILI case definition: Technical Notes


Figure 4



Source of Data: CDC: Flu View Report and CDC: ILINet.

During week 51, 2.6% of visits to sentinel providers were due to ILI. This is a 0.9% point increase from week 51 of the 2012-2013 influenza season.


Figure 5



Source of Data: CDC: ILINet.

The number of ILI patients and rate increased from week 50 to week 51, from 279 to 450, and 10.0 to 16.2 per 100,000 population. The rate is calculated by the number of patients presented with ILI divided by the state population multiplied by 100,000. The estimated state population for 2013 is 2,783,948.


Figure 6



Source of Data: CDC: ILINet.


Providers for the sentinel surveillance are grouped by county, then the percent is calculated by ILI visits and total patient visits. During week 51, Churchill, Douglas, Elko, Eureka, Lander, Pershing, and Washoe counties had high activity; White Pine counties had moderate activity; Humboldt County did not report for week 51 (Figure 7). Overall, Nevada had minimal activity monitored through ILINet (Figure 8).

Figure 7



Source of Data: CDC: ILINet.

Figure 8




Source of Map: CDC: FluView Report.

Pediatric Early Warning Sentinel Surveillance (PEWSS)

Influenza A 2009 H1N1, Human Metapneumovirus, Respiratory Syncytial Virus are detected at sporadic levels, Parainfluenza1 is at a low for week 51.

Figure 9




Source of Data: Southern Nevada Health District: PEWSS.

Influenza Positive Surveillance (NBS and NETSS)

Positive cases of influenza are reported to the state health division for surveillance purposes. Figure 10 and 11 reflect all positive influenza cases reported to the state. Types of influenza testing include commercial rapid diagnostic test (rapid), viral culture, fluorescent antibody, enzyme immunoassay, RT-PCR (PCR), and Immunohistochemistry. The two most common test types in Nevada are Rapid and PCR tests. During week 51, there were 10 H1N1 cases and 253 Influenza A cases. There were 10 positive Influenza B cases. Overall, there were 275 influenza positive tests in Nevada, whereas during the previous season for week 51, there were 100 cases.


Figure 10



Source of Data: OPHIE: NBS and SNHD: NETSS.

Clark County experienced an increase in influenza from week 50 with 65 to 106 influenza cases during week 51. Washoe County experienced an increase in influenza from week 50, from 53 to 104 influenza cases. Carson City, Churchill, Douglas, Elko, Humboldt, Lander, Lyon, Mineral, Pershing, and White Pine counties all had influenza activity during week 51.

Figure 11



Source: OPHIE: NBS and SNHD: NETSS.

Hospitalizations

There have been 63 hospitalizations associated with influenza this season.

Table 2


Influenza Hospitalizations						
Reporting Jurisdiction	Current Week (Week 51)		Cumulative Influenza Season			
	#	%	#	%		
Carson City Health and Human Services	3	15.0	3	4.8		
Rural Health Services	3	15.0	3	4.8		
Southern Nevada Health District	7	35.0	42	66.7		
Washoe County Health District	7	35.0	15	23.8		
State of Nevada	20	100.0	63	100.0		

Source: Reported to Office of Public Health Informatics and Epidemiology from each Jurisdiction.

Pneumonia and Influenza (P&I) Mortality Surveillance

The Pneumonia and Influenza (P&I) mortality percentage is the deaths, where Pneumonia and Influenza is listed as a cause of death, divided by the total deaths in Nevada for each week. There were 29 P&I deaths and 397 total deaths for week 51, as of February 18. The P&I mortality percentage is above the national epidemic threshold at 7.3%, (threshold at 6.9%). Nationally, the P&I mortality is below the epidemic threshold at 6.6%.

Figure 12



Source: OVR: WEVRRS and CDC: FluView.

Appendix

Activity level in figure 3 is based on the following information.

Table 3

Activity Level	ILI Activity*/Outbreaks		Laboratory Data
No Activity	Low	And	
	Not Increased		Isolated lab-confirmed cases †
Sporadic			Or
	Not Increased	And	Lab confirmed outbreak in one institution ‡
	Increased ILI in 1 region**, ILI activity in other regions is not increased	And	Recent (within the past 3 weeks) lab evidence of influenza in region with increased ILI
Local			Or
Local	2 or more institutional outbreaks (ILI or lab confirmed) in 1 region; ILI activity in other regions is not increased	And	Recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions
Dagional	Recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
Regional			Or
	Institutional outbreaks (ILI or lab confirmed) in ≥2 and less than half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions		Recent (within the past 3 weeks) lab confirmed influenza in the state

^{*}ILI activity can be assessed using a variety of data sources including ILINet providers, school/workplace absenteeism and other syndromic surveillance systems that monitor influenza-like illness.

Technical Notes

- Influenza-like illness (ILI): a fever greater than or equal 100°F with cough and/or sore throat
- Percent positive: The number of positive influenza laboratory tests divided by the total number of tests performed.
- Incidence rate is per 100,000 population as estimated by the state demographer.

This report contains information from national and state-level data sources. Influenza surveillance data is collected by a various systems, including:

- Influenza-like Illness Network (ILINet): a sentinel surveillance system in collaboration with the Centers for the Disease Control and Prevention (CDC) where outpatient providers report ILI information weekly.
- National Electronic Telecommunication System for Surveillance (NETSS): a system whereby data is transmits to CDC. Influenza data collected through NETSS does not provide influenza sub-typing information.
- National Electronic Disease Surveillance System (NEDSS): a system for collecting data and monitoring disease trends and outbreaks.
- NEDDS Based System (NBS): an implementation of the NEDSS standards. It provides a secure, accurate, and efficient means of collecting, transmitting, and analyzing public health data.

Citations

- 1. CDC. FluView: A Weekly Influenza Surveillance Report. http://www.cdc.gov/flu/weekly/pastreports.htm.
- 2. Nevada State Demographer's Office. 2003-2012 ASRHO Estimates and Projections. Division of Public and Behavioral Health edition. Vintage 2012.
- 3. OPHIE. DPBH. NBS. 2010-2013. Accessed February 2014.
- 4. Office of Vital Records (OVR). DPBH. Web Enabled Vital Records Registry System (WEVRRS) [unpublished data]. 2012-2013. Accessed February 2014.
- 5. Southern Nevada Health District (SNHD). NETSS/Trisano. 2010-2013. Accessed February 2014.

[†] Lab confirmed case = case confirmed by rapid diagnostic test, antigen detection, culture, or PCR.

[‡] Institution includes nursing home, hospital, prison, school, etc.

^{**}Region: population under surveillance in a defined geographical subdivision of a state. Nevada has 5 regions.

6. SNHD. Pediatric Early Warning Sentinel Surveillance (PEWSS). 2013 PEWSS Reports. Accessed January 2014. http://www.southernnevadahealthdistrict.org/stats-reports/influenza.php.

Comments, suggestions, and requests for further information may be addressed to:

NEVADA INFLUENZA SURVEILLANCE PROGRAM
OFFICE OF PUBLIC HEALTH INFORMATICS AND EPIDEMIOLOGY
4126 TECHNOLOGY WAY, STE 200
CARSON CITY NV 89706
TEL: (775) 684-5897

FAX: (775) 684-5999

Compiled and Written by:

JEN THOMPSON

Recommended Citation:

Division of Public and Behavioral Health. Office of Public Health Informatics and Epidemiology. Influenza Weekly Report, 2012 Week 51 (December 16) through 2013 Week 51 (December 21), Nevada. December 2013 i 51 edition 1.1.

This publication was supported by Cooperative Agreement Number TP000534-02 from the Centers for Disease Control and Prevention and/or Assistant Secretary for Preparedness and Response. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention and/or Assistant Secretary for Preparedness and Response.

