Influenza Weekly Report

2013 Week 06 (February 3 - 9) through 2014 Week 06 (February 2 - 8)

Department of Health and Human Services
Division of Public and Behavioral Health
Office of Public Health Informatics and Epidemiology



Brian Sandoval Governor State of Nevada

Michael J Willden Director Department of Health and Human Services

Richard Whitley, MS Administrator Division of Public and Behavioral Health

Tracey D Green, MD Chief Medical Officer Division of Public and Behavioral Health

Data for the graphs and tables on the following pages are provisional and may be updated as additional information becomes available.


Purpose

The purpose of this report is to provide an overview of and statistics for the influenza season in Nevada for the local public health authorities, sentinel providers and the public.

Influenza-Like Illness Network Surveillance (ILINet)

Respiratory specimens tested for influenza by the World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NRVESS) collaborating laboratories by sub-type. There were 7,562 specimens collected nationally during week 6 that were tested for influenza; of these 1,268 tested positive or the percent positive was 22.7%.


Figure 1



Source of Data: CDC: FluView Weekly Report.

Of the 60 specimens tested for influenza at both the Nevada State Public Health Laboratory and Southern Nevada Public Health Laboratory for sentinel providers, 12 were positive for influenza during week 6 or 20.0%.

Figure 2



Source of Data: CDC: ILINet.

Nevada State Public Health Laboratory (NSPHL) has tested 653 specimens this season with 299 positive from sentinel providers (46.1% positive). Southern Nevada Public Health Laboratory (SNPHL) has reported 42 positive influenza specimens through the Pediatric Early Warning Sentinel Surveillance (PEWSS). Nationally, there have been 189,123 specimens sent to the WHO and NERVSS laboratories with 36,619 positive or 19.4%. The national numbers in table 1 are reflected in figure 1. The state of Nevada data in table 1 is reflected in figure 2.


Table 1

ILINet Surveillance: Influenza Specimens Tested State and Nationally										
	NSPHL SNPHL		State of Nevada (Week 6)		State of Nevada (Season)		National (Week 6)		National (Season)	
			#	%	#	%	#	%	#	%
Specimens Tested	653	240	60		893		7,562		189,123	
Positives to Influenza	301	42	12	20.0	343	38.4	1,268	16.8	36,619	19.4
<u>Influenza A:</u>	299	39	11	91.7	338	98.5	1,154	91.0	35,365	96.6
A(2009 H1N1)	286	38	8	72.7	324	95.9	708	61.4	22,265	63.0
A(Sub-typing not performed)	0	0	0	0.0	0	0.0	401	34.7	12,254	34.7
A(H3)	13	1	3	27.3	14	4.1	45	3.9	845	2.4
<u>Influenza B:</u>	2	3	1	8.3	5	1.5	114	9.0	1,254	3.4

Source of Data: CDC: FluView Report and CDC: ILINet.

For week 6, Nevada reported regional activity to the CDC, along with 19 states (Alabama, Alaska, Colorado, Idaho, Iowa, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Montana, Nebraska, North Dakota, Oregon, South Dakota, Tennessee, Texas, Washington, and Wisconsin). Activity level¹ is derived from data analyzed from Influenza-like Illness (ILI) surveillance (laboratory and sentinel data), and data reported to the state through NBS/NETSS.

Figure 3




Source of Maps: CDC: FluView Report.

Influenza-like Illness (ILI) Network Surveillance has each sentinel providers report the number of patients that meet the ILI case definition² and number of patients that visit the provider weekly, which decreased from 17,512 (week 5) to 16,558 (week 6). The "percentage of visits" is the number of ILI patients divided by the total number of patient visit per week. Nevada's ILI percentage of visits to providers decreased to 2.0% during week 6, and is below the state baseline of 2.3%. Region 9 decreased in ILI to 4.2% from 4.3%, and includes the following states/territories: Arizona, California, Guam, Hawaii, and Nevada. The nation decreased from 3.2% to 3.0% during week 6.

^{1:} Activity level: Appendix Table 4.

^{2:} ILI case definition: Technical Notes


Figure 4



Source of Data: CDC: Flu View Report and CDC: ILINet.

During week 6, 2.0% of visits to sentinel providers were due to ILI. This is a 1.6% point decrease from week 6 of the 2012-2013 influenza season.


Figure 5



Source of Data: CDC: ILINet.

The number of ILI patients and rate decreased from week 5 to week 6, from 394 to 325, and 14.0 to 11.5 per 100,000 population. The rate is calculated by the number of patients presented with ILI divided by the state population multiplied by 100,000. The estimated state population for 2014 is 2,819,321.


Figure 6



Source of Data: CDC: ILINet.


Providers for the sentinel surveillance are grouped by county, then the percent is calculated by ILI visits and total patient visits. During week 6, Elko, and Washoe counties had high activity; Douglas, Eureka, Humboldt, and Lincoln counties did not report for week 6 (Figure 7). Overall, Nevada had low activity monitored through ILINet (Figure 8).

Figure 7



Source of Data: CDC: ILINet.

Figure 8




Source of Map: CDC: FluView Report.

Influenza Positive Surveillance (NBS and NETSS)

Positive cases of influenza are reported to the state health division for surveillance purposes. Figure 10 and 11 reflect all positive influenza cases reported to the state. Types of influenza testing include commercial rapid diagnostic test (rapid), viral culture, fluorescent antibody, enzyme immunoassay, RT-PCR (PCR), and Immunohistochemistry. The two most common test types in Nevada are Rapid and PCR tests. During week 6, there were 18 H1N1 cases and 104 Influenza A cases. There were 24 positive Influenza B cases. Overall, there were 158 influenza positive tests in Nevada, whereas during the previous season for week 6, there were 405 cases.

Figure 9



Source of Data: OPHIE: NBS and SNHD: NETSS.


Table 2

	Reported Influenza Cases by County Jurisdiction and Influenza Type										
Reporting Jurisdiction	Current Week (Week 6)					Cumulative Influenza Season					
	H1N1	Α	В	Unknown	Total	H1N1	A	В	Unknown	Total	
Carson City Health and Human Services	0	6	2	0	8	16	14	15	15	389	
Rural Health Services	4	14	2	0	20	88	26	40	40	381	
Southern Nevada Health District	13	71	24	1	109	1413	100	25	25	1,688	
Washoe County Health District	1	13	7	0	21	482	27	27	27	711	
State of Nevada	18	104	35	1	158	2,466	167	107	107	3,169	

Source: OPHIE: NBS and SNHD: NETSS.

Clark County experienced a decrease in influenza from week 5 with 177 to 109 influenza cases during week 6. Washoe County experienced a decrease in influenza from week 5, from 27 to 21 influenza cases. Carson City, Churchill, Douglas, Elko, and Nye counties all had influenza activity during week 5.

Figure 10



Source: OPHIE: NBS and SNHD: NETSS.

Hospitalizations

There have been 320 hospitalizations associated with influenza this season.

Table 3


Influenza Hospitalizations						
Reporting Jurisdiction	Current We	ek (Week 6)	Cumulative Influenza Season			
	#	%	#	%		
Carson City Health and Human Services	1	6.3	22	6.9		
Rural Health Services	0	0.0	7	2.2		
Southern Nevada Health District	15	93.8	215	67.2		
Washoe County Health District	0	0.0	76	23.8		
State of Nevada	16	100	320	100		

Source: Reported to Office of Public Health Informatics and Epidemiology from each Jurisdiction.

Pneumonia and Influenza (P&I) Mortality Surveillance

The Pneumonia and Influenza (P&I) mortality percentage is the deaths, where Pneumonia and Influenza is listed as a cause of death, divided by the total deaths in Nevada for each week. There were 14 P&I deaths and 259 total deaths for week 6, as of February 18. The P&I mortality percentage is below the national epidemic threshold at 5.4%, (threshold at 7.3%). Nationally, the P&I mortality has surpassed the national epidemic threshold at 8.4%.

Figure 11



Source: OVR: WEVRRS and CDC: FluView.

Appendix

Activity level in figure 3 is based on the following information.

Activity Level	ILI Activity*/Outbreaks		Laboratory Data
No Activity	Low		
	Not Increased		Isolated lab-confirmed cases †
Sporadic			Or
	Not Increased	And	Lab confirmed outbreak in one institution ‡
	Increased ILI in 1 region**, ILI activity in other regions is not increased	And	Recent (within the past 3 weeks) lab evidence of influenza in region with increased ILI
Local			Or
2004.	2 or more institutional outbreaks (ILI or lab confirmed) in 1 region; ILI activity in other regions is not increased	And	Recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions
Dagional	Recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions		Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
Regional			Or
	Institutional outbreaks (ILI or lab confirmed) in ≥2 and less than half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the state

^{*}ILI activity can be assessed using a variety of data sources including ILINet providers, school/workplace absenteeism and other syndromic surveillance systems that monitor influenza-like illness

Nevada has 5 regions.

Technical Notes

- Influenza-like illness (ILI): a fever greater than or equal 100°F with cough and/or sore throat
- Percent positive: The number of positive influenza laboratory tests divided by the total number of tests performed.
- Incidence rate is per 100,000 population as estimated by the state demographer.
- Rapid influenza diagnostic tests Rapid Test.
- Real-time reverse transcription polymerase chain reaction rRT-PCR or PCR.

This report contains information from national and state-level data sources. Influenza surveillance data is collected by a various systems, including:

- Influenza-like Illness Network (ILINet): a sentinel surveillance system in collaboration with the Centers for the Disease Control and Prevention (CDC) where outpatient providers report ILI information weekly.
- National Electronic Telecommunication System for Surveillance (NETSS): a system whereby data is transmits to CDC. Influenza data collected through NETSS does not provide influenza sub-typing information.
- National Electronic Disease Surveillance System (NEDSS): a system for collecting data and monitoring disease trends and outbreaks.
- NEDDS Based System (NBS): an implementation of the NEDSS standards. It provides a secure, accurate, and efficient means of collecting, transmitting, and analyzing public health data.

Citations

- 1. CDC. FluView: A Weekly Influenza Surveillance Report. http://www.cdc.gov/flu/weekly/pastreports.htm.
- 2. Nevada State Demographer's Office. 2003-2012 ASRHO Estimates and Projections. Division of Public and Behavioral Health edition. Vintage 2012.
- 3. OPHIE. DPBH. NBS. 2010-2013. Accessed February 2014.
- 4. Office of Vital Records (OVR). DPBH. Web Enabled Vital Records Registry System (WEVRRS) [unpublished data]. 2012-2013. Accessed February 2014.
- 5. Southern Nevada Health District (SNHD). NETSS/Trisano. 2010-2013. Accessed February 2014.

[†] Lab confirmed case = case confirmed by rapid diagnostic test, antigen detection, culture, or PCR.

[‡] Institution includes nursing home, hospital, prison, school, etc.

^{**}Region: population under surveillance in a defined geographical subdivision of a state.

Comments, suggestions, and requests for further information may be addressed to:

NEVADA INFLUENZA SURVEILLANCE PROGRAM

OFFICE OF PUBLIC HEALTH INFORMATICS AND EPIDEMIOLOGY

4126 TECHNOLOGY WAY, STE 200

CARSON CITY NV 89706

TEL: (775) 684-5897

FAX: (775) 684-5999

Compiled and Written by:

IEN THOMPSON

Recommended Citation:

Division of Public and Behavioral Health. Office of Public Health Informatics and Epidemiology. Influenza Weekly Report, 2013 Week 6 (February 3) through 2014 Week 6 (February 8), Nevada. February 2014 i 6 edition 1.0.

This publication was supported by Cooperative Agreement Number TP000534-02 from the Centers for Disease Control and Prevention and/or Assistant Secretary for Preparedness and Response. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention and/or Assistant Secretary for Preparedness and Response.

