2013 Week 20 (May 12 - 18) through 2014 Week 20 (May 11 - 17)

Department of Health and Human Services Division of Public and Behavioral Health Office of Public Health Informatics and Epidemiology



Brian Sandoval Governor State of Nevada

Michael J Willden Director Department of Health and Human Services

Richard Whitley, MS Administrator Division of Public and Behavioral Health

Tracey D Green, MD Chief Medical Officer Division of Public and Behavioral Health

> June 2014 Edition 1.0

Data for the graphs and tables on the following pages are provisional and may be updated as additional information becomes available.


Purpose

The purpose of this report is to provide an overview of and statistics for the influenza season in Nevada for the local public health authorities, sentinel providers and the public.

Influenza-Like Illness Network Surveillance (ILINet)

Respiratory specimens tested for influenza by the World Health Organization (WHO) and National Respiratory and Enteric Virus Surveillance System (NRVESS) collaborating laboratories by sub-type. There were 3,885 specimens collected nationally during week 20 that were tested for influenza; of these 339 tested positive or the percent positive was 8.7%.


Figure 1



Source of Data: CDC: FluView Weekly Report.

Of the 12 specimens tested for influenza at both the Nevada State Public Health Laboratory and Southern Nevada Public Health Laboratory for sentinel providers, 5 tested positive for influenza during week 20 or 41.7%.

Figure 2



Source of Data: CDC: ILINet.

Nevada State Public Health Laboratory (NSPHL) has tested 809 specimens this season with 328 positive from sentinel providers (40.5% positive). Southern Nevada Public Health Laboratory (SNPHL) has reported 79 positive influenza specimens through the Pediatric Early Warning Sentinel Surveillance (PEWSS). Nationally, there have been 304,291 specimens sent to the WHO and NERVSS laboratories with 52,774 positive or 17.3%. The national numbers in Table 1 are reflected in Figure 1. The state of Nevada data in Table 1 is reflected in Figure 2. The Nevada total includes laboratory test for all Nevada residents including out of state laboratories.


Table 1

ILINet Surveillance: Influenza Specimens Tested State and Nationally										
	NSPHL	SNPHL	State of Nevada (Week 20) # %		State of Nevada (Season)		National (Week 20)		National (Season)	
					#	%	#	%	#	%
Specimens Tested	809	424	12		1,237		3,885		304,291	
Positives to Influenza	328	79	5	41.7	408	33.0	339	8.7	52,744	17.3
Influenza A:	316	43	0	0.0	360	88.2	139	41.0	46,353	87.9
A(2009 H1N1)	301	41	0	0.0	343	95.3	0	0.0	28,322	61.1
A(Sub-typing not performed)	0	0	0	0.0	0	0.0	89	64.0	15,029	32.4
A(H3)	15	2	0	0.0	17	4.7	50	36.0	3,002	6.5
<u>Influenza B:</u>	12	36	5	100	48	11.8	200	59.0	6,390	12.1

Source of Data: CDC: FluView Report and CDC: ILINet.

For week 20, Nevada reported sporadic activity to the CDC, along with 32states/territories and the District of Columbia (Alabama, Alaska, Delaware, Florida, Georgia, Hawaii, Idaho, Illinois, Indiana, Iowa, Kentucky, Louisiana, Michigan, Minnesota, Missouri, Montana, Nebraska, New Mexico, North Dakota, Ohio, Oregon, Pennsylvania, Puerto Rico, South Carolina, South Dakota, Texas, Vermont, Virginia, Washington, West Virginia, Wisconsin, and Wyoming). Activity level¹ is derived from data analyzed from Influenza-like Illness (ILI) surveillance (laboratory and sentinel data), and data reported to the state through NBS/NETSS.

Figure 3




Source of Maps: CDC: FluView Report.

Influenza-like Illness (ILI) Network Surveillance has each sentinel providers report the number of patients that meet the ILI case definition² and number of patients that visit the provider weekly, which decreased to 16,256 (week 20) from 16,398 (week 19). The "percentage of visits" is the number of ILI patients divided by the total number of patient visit per week. Nevada's ILI percentage of visits to providers increased to 0.8% from 0.7% during week 20, and is below the state baseline of 2.3%. Region 9 ILI decreased to 0.9% from 1.4% and includes the following states/territories: Arizona, California, Guam, Hawaii, and Nevada. The national ILI decreased to 1.1% from 1.3% during week 20 and is below the national baseline of 2.0%.

1: Activity level: Appendix Table 4. 2: ILI case definition: Technical Notes


Figure 4



Source of Data: CDC: Flu View Report and CDC: ILINet.

During week 20, 0.8% of visits to sentinel providers were due to ILI. This is a 0.5% point decrease from week 20 of the 2012-2013 influenza season, an influenza season is from week 40 through week 39.


Figure 5



Source of Data: CDC: ILINet.

The number of ILI patients and rate increased from week 19 to week 20, from 110 to 133, and the rate from 3.9 to 4.7 per 100,000 population. The rate is calculated by the number of patients presented with ILI divided by the state population multiplied by 100,000. The estimated state population for 2014 is 2,819,321.


Figure 6



Source of Data: CDC: ILINet.


Providers for the sentinel surveillance are grouped by county, then the percent is calculated by ILI visits and total patient visits. During week 20, Eureka County had high activity; Douglas, Humboldt and Lincoln counties did not report (Figure 7). Overall, Nevada had minimal activity monitored through ILINet (Figure 8).

Figure 7



Source of Data: CDC: ILINet.

Figure 8




Source of Map: CDC: FluView Report.

Influenza Positive Surveillance (NBS and NETSS)

Positive cases of influenza are reported to the state health division for surveillance purposes. Figure 10 and 11 as well as Table 2 reflects all positive influenza cases reported to the state. Types of influenza testing include commercial rapid diagnostic test (rapid), viral culture, fluorescent antibody, enzyme immunoassay, RT-PCR (PCR), and Immunohistochemistry. The two most common test types in Nevada are Rapid and PCR tests. During week 20, there were 27 Influenza A cases. There were 40 positive Influenza B cases. Overall, there were 67 influenza positive tests in Nevada, whereas during the previous season for week 20, there were 15 cases.

Figure 9



Source of Data: OPHIE: NBS and SNHD: NETSS.


Table 2

	Reported Influenza Cases by County Jurisdiction and Influenza Type										
Reporting Jurisdiction	Current Week (Week 20)					Cumulative Influenza Season					
	H1N1	А	В	Unknown	Total	H1N1	А	В	Unknown	Total	
Carson City Health and Human Services	0	4	8	0	12	18	380	41	16	455	
Rural Health Services	0	0	4	0	4	98	249	60	46	453	
Southern Nevada Health District	0	22	25	0	47	177	1,677	444	39	2,336	
Washoe County Health District	0	1	3	0	4	183	513	59	31	786	
State of Nevada	0	27	40	0	67	476	2,819	604	132	4,031	

Source: OPHIE: NBS and SNHD: NETSS.

Clark County experienced a decrease in influenza to 47 from 61 influenza cases during week 20. Washoe County experienced an increase for week 20, to 4 from 2 influenza cases. Carson City, Douglas, and Lyon County had influenza activity during the week.

Figure 10



Source: OPHIE: NBS and SNHD: NETSS.

Hospitalizations

There have been 443 hospitalizations associated with influenza this season (week 40 2013 through week 20).

Table 3


Influenza Hospitalizations						
Reporting Jurisdiction	Current We	ek (Week 20)	Cumulative Influenza Season			
	#	%	#	%		
Carson City Health and Human Services	0	0.0	22	5.0		
Rural Health Services	0	0.0	7	1.6		
Southern Nevada Health District	3	100	313	70.7		
Washoe County Health District	0	0.0	101	22.8		
State of Nevada	3	100	443	100		

Source: Reported to Office of Public Health Informatics and Epidemiology from each Jurisdiction.

Pneumonia and Influenza (P&I) Mortality Surveillance

The Pneumonia and Influenza (P&I) mortality percentage is the deaths, where Pneumonia and Influenza is listed as a cause of death, divided by the total deaths in Nevada for each week. There were 23 P&I deaths and 342 total deaths for week 20, as of June 2nd. The P&I mortality percentage is above the national epidemic threshold at 6.7%, (threshold at 6.9%). Nationally, the P&I mortality is below the national epidemic threshold at 5.8%.

Figure 11



Source: OVR: WEVRRS and CDC: FluView.

Appendix

Activity level in figure 3 is based on the following information.

Activity Level	ILI Activity*/Outbreaks		Laboratory Data
No Activity	Low	And	
	Not Increased		Isolated lab-confirmed cases †
Sporadic			Or
	Not Increased	And	Lab confirmed outbreak in one institution ‡
	Increased ILI in 1 region**, ILI activity in other regions is not increased		Recent (within the past 3 weeks) lab evidence of influenza in region with increased ILI
Local			Or
2000	2 or more institutional outbreaks (ILI or lab confirmed) in 1 region; ILI activity in other regions is not increased		Recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions
Decional	Recent (within the past 3 weeks) lab evidence of influenza in region with the outbreaks; virus activity is no greater than sporadic in other regions		Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
Regional			Or
	Institutional outbreaks (ILI or lab confirmed) in ≥2 and less than half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the affected regions
Widespread	Increased ILI and/or institutional outbreaks (ILI or lab confirmed) in at least half of the regions	And	Recent (within the past 3 weeks) lab confirmed influenza in the state

*ILI activity can be assessed using a variety of data sources including ILINet providers, school/workplace absenteeism and other syndromic surveillance systems that monitor influenzalike illness.

† Lab confirmed case = case confirmed by rapid diagnostic test, antigen detection, culture, or PCR.

‡ Institution includes nursing home, hospital, prison, school, etc.

**Region: population under surveillance in a defined geographical subdivision of a state. Nevada has 5 regions.

Technical Notes

- Influenza-like illness (ILI): a fever greater than or equal 100°F with cough and/or sore throat
- Percent positive: The number of positive influenza laboratory tests divided by the total number of tests performed.
- Incidence rate is per 100,000 population as estimated by the state demographer.

This report contains information from national and state-level data sources. Influenza surveillance data is collected by a various systems, including:

- Influenza-like Illness Network (ILINet): a sentinel surveillance system in collaboration with the Centers for the Disease Control and Prevention (CDC) where outpatient providers report ILI information weekly.
- National Electronic Telecommunication System for Surveillance (NETSS): a system whereby data is transmits to CDC. Influenza data collected through NETSS does not provide influenza sub-typing information.
- National Electronic Disease Surveillance System (NEDSS): a system for collecting data and monitoring disease trends and outbreaks.
- NEDDS Based System (NBS): an implementation of the NEDSS standards. It provides a secure, accurate, and efficient means of collecting, transmitting, and analyzing public health data.

Citations

- 1. CDC. FluView: A Weekly Influenza Surveillance Report. <u>http://www.cdc.gov/flu/weekly/pastreports.htm</u>.
- 2. Nevada State Demographer's Office. 2003-2014 ASRHO Estimates and Projections. Division of Public and Behavioral Health edition. Vintage 2012.
- 3. OPHIE. DPBH. NBS. 2010-2014. Accessed June 2014.
- 4. Office of Vital Records (OVR). DPBH. Web Enabled Vital Records Registry System (WEVRRS) [unpublished data]. 2012-2014. Accessed June 2014.
- 5. Southern Nevada Health District (SNHD). NETSS/Trisano. 2010-2014. Accessed June 2014.

Comments, suggestions, and requests for further information may be addressed to:

NEVADA INFLUENZA SURVEILLANCE PROGRAM OFFICE OF PUBLIC HEALTH INFORMATICS AND EPIDEMIOLOGY 4126 TECHNOLOGY WAY, STE 200 CARSON CITY NV 89706 TEL: (775) 684-5897 FAX: (775) 684-5999

Compiled and Written by:

JEN THOMPSON

Recommended Citation:

Division of Public and Behavioral Health. Office of Public Health Informatics and Epidemiology. Influenza Weekly Report, 2013 Week 20 (May 12) through 2014 Week 20 (May 17), Nevada. June 2014 i 20 edition 1.0.

This publication was supported by Cooperative Agreement Number TP000534-02 from the Centers for Disease Control and Prevention and/or Assistant Secretary for Preparedness and Response. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention and/or Assistant Secretary for Preparedness and Response.

