

Office of Analytics
Department of Health & Human Services

Treatment Episode Data Sets (TEDS) – Discussion Points

There were 7,082 admissions to substance use treatment facilities at the end of the 3rd quarter.

Type of treatment service/treatment setting

Admissions to Ambulatory - Non-Intensive Outpatient Treatment Services decreased from about 54% to 40% of total services. Admissions to Detoxification - Free-standing Residential Services increased from about 17% to 37% of total services.

Primary Source of Referral

In the 3rd quarter, 2,562 admissions were self-referred compared to 552 in the 2nd quarter.

In the 3rd quarter, 2,163 admissions were referred by the Courts/Criminal Justice System as compared to 1,145 in the 2nd quarter.

Gender

In the 3rd quarter, 4,441 males and 2,640 females sought help as compared to 1,702 males and 1,083 females in the 2nd.

Race

In the 3rd quarter, around 59% of admissions were white and 15% were black; roughly 64% were white & 11% were black in the 2nd.

Hispanic or Latino origin

In the 3rd quarter, roughly 63% were not of Hispanic origin as compared to 64% in the 2nd quarter.

Substance Use (primary)

Methamphetamine constituted about 16% of total drug use in the 3rd quarter as compared to 21% in the 2nd.

Alcohol constituted roughly 15% of total drug use in the 3rd quarter as compared to 19% in the 2nd.

Route of Administration (primary)

In the 3rd quarter, smoking as a route of administration was about 21% as compared to 28% in the 2nd.

In the 3rd quarter, oral route of administration constituted about 16% as compared to 21% in the 2nd.

Frequency of Use (primary)

Daily use was roughly 29% in the 3rd quarter and 36% in the 2nd quarter.

Co-occurring Mental and Substance Use Disorders

In the 3rd quarter, about 21% of admissions had a co-occurring disorder and 28% exhibited the same in the 2nd.

Days Waiting to Enter Treatment

In the 3rd quarter, about 74% of admissions waited for an average of 0-5 days as compared to 46% in the 2nd.

Substance Use Diagnosis

Opioid dependence was diagnosed in about 9% of admissions in the 2nd & 3rd quarters.