



## Influenza Surveillance Report – 2019-2020 Season – Week 11

### Data from March 8 – 14, 2020

#### Introduction

The purpose of this report is to provide ongoing description and assessment of the activity and types of circulating influenza viruses, and to assess morbidity, hospitalization and mortality related to influenza. It is meant to provide healthcare providers and facilities, public health professionals, policy makers, the media and the public with a general understanding of the severity and burden of the current flu season on a weekly basis in Nevada and nationwide. Data from several surveillance programs analyzed in this report is provisional and may change as additional information becomes available.

If you have questions or comments about this report, are interested in having your medical facility join the sentinel provider program, or have any questions about your facility’s participation or reporting, please contact Max Wegener, MPH at [mwegener@health.nv.gov](mailto:mwegener@health.nv.gov).

**Influenza activity in the State of Nevada is presently widespread:** outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least half the regions of the state with recent laboratory evidence of influenza in the state.

Table 1:

Week 11 Summary					
	ILI Current Activity	ILI Activity Baseline	Influenza -related Hospitalization	Influenza -related Mortality	Pneumonia and Influenza Mortality
Nevada	3.09%	1.36%	21(0.7 per 100,000)	0/445 (0%)	27/445 (6.07%)*
Region 9	4.02%	2.40%	pending	63/6947 (0.91%)*	515/6947 (7.41%)*
National	5.65%	2.20%	1 per 100,000	491/47655 (1.03%)*	3203/47655 (6.72%)*

\*CDC data based on cause of death listed in vital records

#### Local Health Authority (LHA) reports

Weekly influenza reports from the three LHAs are available on the respective websites:

- Southern Nevada Health District: <https://www.southernnevadahealthdistrict.org/stats-reports/influenza-surveillance.php>
- Washoe County Health District: [https://www.washoecounty.us/health/programs-and-services/communicable-diseases-and-epidemiology/statistics\\_surveillance\\_reports/influenza-surveillance/index.php](https://www.washoecounty.us/health/programs-and-services/communicable-diseases-and-epidemiology/statistics_surveillance_reports/influenza-surveillance/index.php)
- Carson City Health & Human Services: Western NV Regional Influenza Report: <http://gethealthycarsoncity.org/seasonalflu/>

## Sentinel Provider Program Description

The sentinel provider program is a partnership between clinicians, healthcare facilities, local health authorities (LHA), the Nevada Division of Public and Behavioral Health, and the Centers for Disease Control and Prevention (CDC). Sentinel providers voluntarily submit a weekly report to the CDC of the number of patients seen at their facility with influenza-like illness (ILI) by age group as well as the total number of patients seen for any reason. ILI is defined as fever ( $\geq 100^{\circ}\text{F}$ ,  $37.8^{\circ}\text{C}$ ) in the presence of cough and/or sore throat without a known cause other than influenza. Sentinel providers may also submit nasal, throat, and/or nasopharyngeal swabs for selected patients to the Nevada State Public Health Laboratory (NSPHL) for viral testing and subtyping at no cost to the patient or provider.

## Sentinel Provider Influenza-Like Illness (ILI) Activity:

**Figure 1** shows the percent of ILI patients by age group for week 11. Those age 0-4 represented 33% of all reported ILI cases in Nevada. 29% of cases were ages 5-24, 21% ages 25-49, 8% ages 50-64, and 9% ages 65 and older.

In week 11, 9,871 patient visits were reported by sentinel providers in Nevada, of which 305 met criteria for ILI, representing 3.1% of the sample. ILI activity was above the Nevada baseline of 1.4%. **Figure 2** shows the percent of reported visits statewide for which the patient met clinical criteria for ILI. The current influenza season (2019 week 40 – 2020 week 20), in bold, is overlaid with the prior four seasons.

For week 11, 4.0% of patients reported in Region 9 (AZ, CA, HI, NV, and US Pacific Islands) and 5.6% of patients reported nationally met criteria for ILI. The regional activity level is greater than the regional baseline of 2.4% and the national activity level is greater than the national baseline of 2.2%.

**Figure 3** displays a comparison of the percent of visits which met ILI criteria for Nevada, Region Nine, and nationally.

Figure 1:

2020 Week 11  
Percent of ILI patients by age




Figure 2:





Figure 3:



### Sentinel Providers Virologic Testing

The Nevada State Public Health Laboratory (NSPHL) and other laboratories provide influenza virus testing and subtyping for specimens submitted by sentinel providers. For week 11, 19 specimens were positive of 96 submitted (19.8%). From week 40 to date, 880 specimens were positive of 1,826 submitted (48%). **Figure 4** shows the number of laboratory-confirmed influenza cases by subtype expressed as a percentage of all laboratory-confirmed specimens tested. Of the 880 positive specimens to date, 34 were typed as influenza A (H3N2), 263 as A (2009 H1N1), 223 as A (subtyping not performed), 131 as B (Victoria), and 229 as B (subtyping not performed). **Table 2** shows the number of sentinel site specimens tested by laboratory this season and the number and percent positive for influenza of any type.

Figure 4:




Table 2:

Lab	# of tests performed	# positive	% positive
Nevada State Public Health Lab (NSPHL)	353	259	73%
Southern Nevada Public Health Lab (SNPHL)	265	170	64%
All other labs	1,208	451	37%
Total	1,826	880	48%

## Influenza Hospitalizations

LHAs investigate and report to DPBH Influenza-associated hospitalizations. **Figure 5** shows the number of patients hospitalized with influenza by jurisdiction. In week 11, Southern Nevada Health District reports 21; Washoe County Health District, Carson City Health and Human Services, and Rural Health Services report none. From week 40 to date, 1,609 total hospitalizations have been reported statewide. **Figure 6** shows the number of hospitalized patients by influenza type, if reported.

Figure 5:






Figure 6:



## Influenza Deaths

Influenza-associated deaths are deaths from a clinically-compatible illness that was confirmed to be influenza by an appropriate laboratory or rapid diagnostic test with no period of complete recovery between illness and death. LHAs investigate all influenza deaths and typically review medical records retroactively up to 30 days from the date of death for an influenza diagnosis. **Figure 7** shows the number of influenza deaths by region for this flu season. No deaths were reported in week 11. There have been 46 influenza-associated deaths reported statewide since week 40.

**Figure 7:**



**Syndromic Surveillance**

Syndromic surveillance uses near real-time, pre-diagnostic health data to analyze disease incidence. It may support the identification and characterization of outbreaks as supplemental data or as an early indicator of a possible outbreak. DPBH uses the National Syndromic Surveillance System (NSSP) Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE), a CDC web application, to collect these data from hospitals and urgent care facilities within the state. Chief complaint is used for immediate analysis; discharge diagnosis is used as it becomes available.

**Syndromic Surveillance ILI Activity**

**Figure 8** shows the number of visits with ILI for emergency, inpatient, and outpatient settings. For week 11 there were 1,335 emergency visits, 47 hospital admissions, and 107 outpatient visits reported. Emergency department visits increased by 32% from 1,012 the previous week. **Figure 9** shows the number of emergency visits with ILI by week over five years; **figure 10** shows the number of inpatient visits with ILI by week over five years. **Figure 11** shows the percent of all visits with ILI by age group. For week 11, 22% of visits were for ages 0-4, 26% for ages 5-24, 33% for ages 25-49, 11% for ages 50-64, and 8% for ages 65 and up.

**Figure 8:**






Figure 9:




Figure 10:





Figure 11:



## Pneumonia and Influenza (P&I) Mortality Surveillance


Death certificate data are used to calculate pneumonia and influenza deaths. The Division of Public and Behavioral Health is presently evaluating its data extraction methodology and will report P&I deaths in the future from internal data.

The CDC makes P&I death information available in its FluView Interactive GIS application. According to data from the CDC, Nevada's P&I mortality is 6.07% of all deaths reported (27 out of 445) for the most recent week. Region 9's P&I mortality is 7.41% of all deaths reported, which is above the baseline of 7.3%; nationally 6.72% of all deaths are due to P&I, which is below the baseline of 6.92%. Region 9's influenza-related mortality is 0.91% (63 out of 6,947) and nationally 1.03% of all deaths are influenza-related (491 out of 47,655).

## Respiratory syncytial virus (RSV)

From week 40 through week 11, 2,519 Respiratory Syncytial Virus (RSV) cases have been reported. **Figure 12** shows the number of reported RSV cases for the current season compared with the number reported in the past four seasons. Due to a lag in reporting the case count for recent weeks is expected to increase.

Figure 12:



## References

Figures 1, 2, and 3, and Table 1 are derived from ILINet sentinel surveillance data submitted by sentinel providers directly to the CDC.

Table 1 also uses data from CDC's FluView Interactive GIS application.

Figure 4 and Table 2 use ILINet laboratory surveillance data.

Figures 5, 6, and 7 are compiled from data collected by local health authorities and abstracted from medical records.

Figures 8, 9, 10, and 11 are populated from the National Syndromic Surveillance System (NSSP) Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE).

Figure 12 is generated from data submitted to Nevada's NBS/NETSS reporting systems.