EPIDEMIOLOGIC INVESTIGATION SUMMARY

Norovirus Outbreak Among Residents and Staff of an Assisted Living Facility in Washoe County, Nevada, 2017

Department of Health and Human Services
Division of Public and Behavioral Health
Office of Public Health Informatics and Epidemiology

November 2017 Edition 1.0 2017 volume, issue 31

PURPOSE

The purpose of this newsletter is to provide the scientific community, decision makers, healthcare providers, and the public a summary of the outbreak investigations conducted by the Division of Public and Behavioral Health.

BACKGROUND

On November 6, 2017, the Division of Public and Behavioral Health (DPBH), Office of Public Health Informatics and Epidemiology (OPHIE) was informed of a gastrointestinal (GI) illness among residents of Facility "A." The outbreak or increase in illness was first identified by staff of the facility on November 4, 2017. Initial symptomology of the ill residents included vomiting and diarrhea. The outbreak investigation began on November 6, 2017.

CASE DEFINITIONS

Clinical criteria An illness with gastroenteritis that presents with the following symptoms: vomiting, diarrhea, abdominal cramps or stomach ache, nausea, with or without fever between November 3, 2017 to November 11, 2017.

Epidemiological criteria Any residents or staff members associated with Facility "A" identified through investigations.

Laboratory criteria Any laboratory confirmation by PCR or other method from a human specimen for an enteric etiology.

Case classification

A **confirmed case** A case meeting clinical, epidemiological, and laboratory criteria.


A **probable case** A case meeting the following clinical criteria and epidemiological criteria.


Vomiting OR Diarrhea OR Abdominal cramping (stomach ache) + fever* OR Nausea + fever*
*fever refers to self-reported fever or measured elevated temp.

Epidemiology

Onset Date

The peak illness onset date was November 6, 2017.





Epidemiology Summary

A total of 12 cases met the confirmed and probable case definition (two lab-confirmed and ten probable). There were no deaths associated with this outbreak and there were 3 hospitalizations (all residents). The resident attack rate was 38.5% (n=26) and the staff attack rate was 10.5% (n=19). The overall attack rate was 26.7%. *Age-* The median age was 72 (range: 19 – 94 years).

Age	n	Total N	%
10-19 years	1	12	8.3%
20-49 years	1	12	8.3%
50-74 years	2	12	16.7%
> = 75	8	12	66.7%

Sex- Male n=2 (16.7%), Female n=10(83.3%)

Incubation period- The incubation period for norovirus is 12-48 hours¹.

Duration of illness- The average duration of illness was approximately two days (range one – four days).

Summary of Symptoms-

Symptoms	n	Total N	%
Diarrhea	8	12	67%
Fever	1	12	8.3%
Nausea	1	12	8.3%
Vomiting	6	12	50%

Laboratory

There was a total of two specimens tested, both of which were positive for norovirus genogroup II. One of the positive specimens was also positive for Clostridium *difficile*.

Data Sources

Residents who reported complaints consistent with GI illness. (line listing form)

Staff who called in with complaints consistent with GI illness. (line listing form)

CONCLUSIONS

The latest onset date occurred on November 11, 2017. The facility completed two incubation periods with no new cases, therefore the outbreak investigation was closed on November 21, 2017.

Mitigation

After lab results confirmed that the cause of the outbreak was norovirus, which has an incubation period of 12-48 hours¹, DPBH reiterated the importance of continued outbreak control measures to interrupt further transmission. The facility continued their own mitigation efforts as well.

RECOMMENDATIONS

To prevent norovirus outbreaks in healthcare settings, the following public health measures are recommended:

- Follow hand-hygiene guidelines and careful washing of hands with soap and water after contact with patients with norovirus infection.
- Use gowns and gloves when in contact with or caring for patients who are symptomatic with norovirus.
- Routinely clean and disinfect high touch patient surfaces and equipment with an Environmental Protection Agency-approved product with a label claim for norovirus.

- After vomiting or having diarrhea, immediately clean and disinfect contaminated surfaces using a bleachbased household cleaner, which is effective against norovirus, as directed on the product label. If no such cleaning product is available, you can mix a solution with ½ cup of bleach to one gallon of water.
- Remove and wash contaminated clothing and linens.
- Exclude healthcare workers who have symptoms consistent with norovirus from work for 48 hours after the resolution of symptoms.
- During outbreaks, place patients with norovirus gastroenteritis on Contact Precautions for a minimum of 48 hours after the resolution of symptoms to prevent further exposure to susceptible patients.

REFERENCES

1. https://www.cdc.gov/hai/pdfs/norovirus/229110-anorocasefactsheet508.pdf

For additional information regarding this publication, contact:

Office of Public Health Informatics and Epidemiology 4126 Technology Way, Ste 200 Carson City NV 89706 Email: outbreak@health.nv.gov Tel: (775) 684-5911



Brian Sandoval Governor State of Nevada

Richard Whitley, MS
Director
Department of Health and Human Services

Julie Kotchevar, PhD Interim Administrator Division of Public and Behavioral Health

Dr. Leon Ravin, MD
Chief Medical Officer (Acting)
Division of Public and Behavioral Health



RECOMMENDED CITATION

Division of Public and Behavioral Health. Office of Public Health Informatics and Epidemiology. Epidemiologic Investigation Summary, *Norovirus Outbreak among Residents and Staff of an Assisted Living Facility in Washoe County, Nevada, 2017.* v 2017. i 31. e 1.0. November 2017.

ACKNOWLEDGEMENTS

Thank you to all persons who contributed to this publication:

Jessica Conner, MPH; Brian Parrish, MPH; Kimisha Causey, MPH; Adrian Forero, BS; Chidinma Njoku BS; Judy Dumonte; Sandi Larson, MPH; Melissa Peek-Bullock; Ihsan Azzam, MD, MPH.

This report was produced by the Office of Public Health Informatics and Epidemiology of the Division of Public and Behavioral Health with funding from budget account 3219.